

Ciprofloxacin

You are being given this medicine because you may have been exposed to Anthrax. You will need to take this medicine for 60 days. You will be told how to get more medicine if it is needed.

Dosing Information

- Take the medicine every 12 hours or as directed.
- **Do not take antacids** such as Tums, Rolaids or milk of magnesia 2 hours before taking this medicine. Also, wait 2 hours after taking the medicine before you take any antacids. Antacids can lessen how well the medicine works.
- Take the medicine with a full glass of water.
- If you miss a dose, take the missed dose right away. If it is almost time to take your next dose, wait until then to take your medicine and skip the missed dose. Do not take two doses at the same time.
- You must keep taking your medicine for the full 60 days. You may become ill if you do not finish all your medicine.

Side Effects

- This medicine can cause upset stomach, vomiting, diarrhea, headaches, dizziness, or restlessness, but **you need to keep taking this medicine if any of these occur.**
- **Call your doctor right away** if you have:
 - trouble breathing
 - swelling of the face, lips, tongue or throat
 - if you have a skin rash with mouth blisters
 - shortness of breath

Interactions

- If you take warfarin (Coumadin), phenytoin (Dilantin), digoxin, theophylline, Requip, cyclosporine or probenecid, **call 211 or call your doctor and tell them that you are taking Ciprofloxacin.** These medicines can cause problems if taken while taking Ciprofloxacin.
- Do not use any herbs, especially St. John's Wort and Dong Quai while taking this medicine.
- If you are on kidney dialysis, tell your center that you are taking Ciprofloxacin.

Ciprofloxacin

Waxa lagu siinayaa dawadan sababta oo ah waxa ku soo gaadhay Kud. Waxa aad u baahan tahay inaad qaadatid dawadan 60 maalmood. Waxa lagu sheegi doonaa sida aad ku heli kartid dawo dheeraad ah, haddii loo baahdo.

Warka Qadarka Qaadashada

- Qaado dawada 12 saacadood kasta ama u qaado sida lagu faro.
- **Ha qaadan dawooyinka lidka asiidhka ama aashitada** sida Tums, Roloids ama milk (caano) of magnesia 2 saacadood ka hor qaadashada dawadan. Sidoo kale, marka aad qaadatid dawada sug 2 saacadood, ka hor inta aadan qaadan wax ah lidka asiidhka ama aashitada. Dawooyinka lidka asiidhku waxay yareyn karaan shaqeynta dawadan.
- Dawada ku qaado koob buuxa oo ah biyo.
- Haddii aad gaftid hal qaadasho, isla markiiba qaado qadarka qaadashado ee aad gaftay. Haddii uu soo dhow yahay wakhtigii qaadashada kale, sug ka dibna qaado dawadaada oo ka bood qadarka aad gaftay. Isku wakhti ha isku qaadan laba qadar qaadasho.
- Waa inaad sii wadid qaadashada dawadaada mudada buuxda ee ah 60 maalmood. Waad xanuunsan kartaa haddii aada qaadan dhammaan dawadaada.

Saameynnada

- Dawadani waxay sababi kartaa qas calool, matag, shuban, madax-xanuun, dawakh (wareer), ama deganaansho la'aan, laakiin **waxa la rabaa inaad sii wadid qaadashada dawadan haddii mid ka mid ah kuwaas ay dhacaan.**
- **Isla markiiba wac dhakhtarkaaga** haddii aad yeelatid:
 - neefsashada oo dhib kugu ah
 - barar ah wajiga, dibnaha, carrabka ama cunaha
 - haddii aad leedahay finan maqaarka ah ayna jiraan finan ama boogo afka ah
 - neefsashada oo gaaban

Falgalka Dawooyinka

- Haddii aad qaadatid warfarin (Coumadin), phenytoin (Dilantin), digoxin, theophylline, Requip, cyclosporine ama probenecid, **wac 211 ama wac dhakhtarkaaga oo u sheeg inaad qaadatid Ciprofloxacin.** Dawooyinkani waxay sababi karaan dhibaatooyin haddii lala qaato Ciprofloxacin.
- Ha isticmaalin wax ah geedo ama dhir, gaar ahaan St. John's Wort iyo Dong Quai mudada aad qaadanaysid dawadan.
- Haddii lagu sameeyo ka-sifeyn-kaadi (dialysis), u sheeg xaruntaada inaad qaadatid Ciprofloxacin.