


# Ciprofloxacin Suspension for Babies

## How to give Ciprofloxacin Suspension To Your Child


- Mix the medicine as shown to the right.
- Shake the medicine for about 15 seconds before measuring out the next dose.
- After it is mixed, keep the medicine in the refrigerator. It may be kept covered in the refrigerator for 14 days.
- **Give two doses each day. Give one in the morning and one in the evening.**
- Measure the amount of liquid medicine for your child's dose, as marked below.
- To make it taste better, the suspension can be mixed with a small amount of food or juice. **Be sure the child takes all this mixture to get the full dose of medicine.**

Child's weight	Give this much	How often each day
<b>7-12 pounds</b> (3-5 kg)	<b>1 ml</b>	Two times
<b>13-22 pounds</b> (6-10 kg)	<b>2 ml</b>	Two times
<b>23-33 pounds</b> (11-15 kg)	<b>3 ml</b>	Two times
<b>34-44 pounds</b> (16-20 kg)	<b>4 ml</b>	Two times


## Mixing Instruction:




1. The small bottle contains the medicine. **Do not chew the medicine.** Mix the medicine with the liquid in the larger bottle.



2. Open both bottles. Press down on the cap as you twist it to the left.



3. Pour the medicine into the larger bottle of liquid. **Do not add water or any other liquid.** Be sure you get all the medicine out of the small bottle.




4. Close the large bottle. Be sure the cap is on tight. Shake the bottle well for about 15 seconds. The medicine is ready for use.

## 如何让孩子服用 Ciprofloxacin悬浮液


- 按右图所示，混合药物。
- 量出所服药量之前，摇匀大约15秒。
- 混合后，放入冰箱内保存。加盖后，可在冰箱内保存14天。
- **每天服用两次。早晚各一次。**
- 按下表所示，量出孩子要服的药。
- 为了改善药物口味，可与少量食物或果汁混合后让孩子服用。**务必要让孩子将混合物全部服下，以确保全剂量服药。**

儿童体重	服用量	每日服用次数
<b>3-5公斤</b> (7-12磅)	<b>1 ml</b>	两次
<b>6-10公斤</b> (13-22磅)	<b>2 ml</b>	两次
<b>11-15公斤</b> (23-33磅)	<b>3 ml</b>	两次
<b>16-20公斤</b> (34-44磅)	<b>4 ml</b>	两次


## 混合方法




1. 小瓶内是药物。**切勿嚼服。**将药物与大瓶内的液体混合。



2. 打开两个瓶子，方法是下压瓶盖，向左转动。



3. 将药物倒入装液体的大瓶内。**切勿加水或任何其它液体。**务必倒空小瓶内的药物。



4. 盖上大瓶。瓶盖一定要拧紧。摇动药瓶约15秒后，即可服用。