



Giving Doxycycline Pills to Your Child



- The amount of medicine depends on the child’s weight.
- For a child who weighs less than 55 pounds (25kg), you should use the Doxycycline Suspension for your child, not the pills. Please refer to that page for directions.
- **Give two doses each day. Give one in the morning and one in the evening.** Give the amount marked below for your child.
- To make it taste better, the medicine can be crushed and mixed with a **small amount** of food or juice. **Be sure the child takes all this mixture to get the full dose of medicine.**

Child’s weight	Morning Dose Amount	Evening Dose Amount
Under 55 pounds (under 25 kg)	See the page about Doxycycline Suspension for medicine directions	
55-80 pounds (25-36 kg)	½ pill	½ pill
81-98 pounds (37-44 kg)	1 pill	½ pill
99 pounds or more (45kg or more)	1 pill	1 pill



兒童服用Doxycycline藥片



- 藥物劑量依兒童體重而定。
- 體重不足25公斤（55磅），應使用Doxycycline懸浮液，勿使用藥片。請參閱有關說明。
- 每天服用兩次。早晚各一次。按下表的藥量給孩子服藥。
- 爲了改善藥物的味道，可將藥片弄碎後與少量食物或果汁混合後服用。務必要讓孩子將混合物全部服下，以確保全劑量服藥。

兒童體重	早晨劑量	晚上劑量
25公斤以下（55磅以下）	參閱Doxycycline懸浮液服用說明	
25-36公斤（55-80磅）	½片	½片
37-44公斤（81-98磅）	1片	½片
45公斤或以上 （99磅或以上）	1片	1片

Directions for Anthrax Prevention for Children Over 54 pounds
with Doxycycline 100mg tablets. Traditional Chinese

Adapted by the Communicable Disease Prevention Team of Columbus Public Health from protocols developed by the Ohio Dept. of Health 3/08