



Giving Doxycycline Pills to Your Child



- The amount of medicine depends on the child’s weight.
- For a child who weighs less than 55 pounds (25kg), you should use the Doxycycline Suspension for your child, not the pills. Please refer to that page for directions.
- **Give two doses each day. Give one in the morning and one in the evening.** Give the amount marked below for your child.
- To make it taste better, the medicine can be crushed and mixed with a **small amount** of food or juice. **Be sure the child takes all this mixture to get the full dose of medicine.**

Child’s weight	Morning Dose Amount	Evening Dose Amount
Under 55 pounds (under 25 kg)	See the page about Doxycycline Suspension for medicine directions	
55-80 pounds (25-36 kg)	½ pill	½ pill
81-98 pounds (37-44 kg)	1 pill	½ pill
99 pounds or more (45kg or more)	1 pill	1 pill

Sida Kiniinka Doxycycline Loo Siiyo Ilmahaaga

- Qadarka dawadu waxa uu ku xidhan yahay miisaanka ilmahaaga.
- Marka ilmaha miisaankiisu ka yar yahay 25 kg (55 rodol), waa inaad ilmahaaga siisid Doxycycline Suspension oo ah hoore, oo aadan siinin dawada oo kiniin ah. Fadlan bogga hore ka eeg farriimo.
- **Maalin kasta sii laba qadar qaadasho. Hal sii subixii, halna sii fiidkii.** Qadarka hoos loogu calaamadeeyay sii ilmahaaga.
- Si dhadhanku u fiicnaado, waxa dawada hooraha ah lagu qasi karaa **qadar yar** oo ah cunto ama miir khudaar. **Hubso in ilmuhu qaato dhammaan isku-jirka si uu u helo qadar buuxa oo ah dawada.**

Miisaanka ilmaha	Qadarka Qaadashada ee Subixii	Qadarka Qaadashada ee Fiidkii
Ka yar 25 kg (ka yar 55 rodol)	Eeg bogga ku saabsan Doxycycline Suspension si aad u ogaatid farriimaha dawada	
25-36 kg (55-80 rodol)	½ kiniin	½ kiniin
37-44 kg (81-98 rodol)	1 kiniin	½ kiniin
45 kg ama ka badan (99 rodol ama ka badan)	1 kiniin	1 kiniin