


Doxycycline Suspension for Babies


How to give Doxycycline Suspension To Your Child

- Mix the medicine as shown to the right.
- Shake the medicine for about 15 seconds before measuring out the next dose.
- After it is mixed, keep the medicine in the refrigerator. It may be kept covered in the refrigerator for 14 days.
- Measure the amount of liquid medicine for your child's dose, as marked below.
- Give 1 or 2 doses each day. Give the medicine in the morning. If your child is to have 2 doses each day, give the medicine in the evening also. Follow the schedule marked below for your child.
- To make it taste better, the suspension can be mixed with a small amount of food or juice. **Be sure the child takes all this mixture to get the full dose of medicine.**


Mixing Instructions




1. Open the bottle. Press down on the cap as you twist it to the left.



2. Measure 1.5 ounces of water in the medicine cup you were given. (45ml = 1.5 ounces)



3. Add the water to the medicine powder.




4. Close the bottle. Be sure the cap is on tight. Shake the bottle well for about 15 seconds. The medicine is ready for use.

Child's weight	Give this much liquid	How often each day
Under 12.5 pounds (<6 kg)	5 ml	One time
12.5-27 pounds (6-12 kg)	10 ml	One time
28-54 pounds (13-24 kg)	15 ml	One time


Sida Hooraha Doxycycline Suspension Loo Siyo Ilmahaaga

- Dawada u qas sida lagu muujiyay midigta.
- Rux dawada qiyaas ah 15 sekan (il-biriqsi) ka hor cabirista qadarka qaadashada ee ku xiga.
- Ka dib marka la qaso, dawada ku hay qaboojiyaha. Iyada oo daboolan, waxa lagu hayn karaa qaboojiyaha mudo ah 14 maalmood.
- Sida hoos lagu calaamadeeyay, u cabir qadarka qaadashada ilmahaaga ee dawada hooraha ah.
- Sii 1 ama 2 qadar qaadasho maalin kasta. Sii dawada subaxii. Haddii la rabo in ilmahaaga la siiyo 2 qadar qaadasho maalin kasta, markale sii dawada fiidkii. Raac jadwalka hoos loogu calaamadeeyay ilmahaaga.
- Si dhadhankeeda loo fiicneeyo, waxa hooraha lagu qasi karaa qadar yar oo ah cunto ama miir khudaar. **Hubso in ilmuhu qaato dhammaan isku-darka si uu u helo qadar buuxa oo ah dawada.**


Farriimaha Qasidda




1. Fur dhalada. Hoos u riix furka adiga oo u wareejinaya bidix.



2. Koobka dawada ee lagu siiyay ku cabir 1.5 wiqiyadood (ounces) oo ah biyo. (45 ml = 1.5 wiqiyadood)



3. Biyaha ku dar dawada budada ah.



4. Xidh dhalada. Hubso in furku ku adag yahay. Si fiican u rux dhalada qiyaas ah 15 sekan (il-biriqsi). Dawadu waa u diyaar in la isticmaalo.

Miisaanka ilmaha	Sii qadarkan ah hooraha	Intee jeer maalin kasta
Ka yar 6 kg (<12.5 rodol)	5 ml	Hal wakhti
6-12 kg (12.5-27 rodol)	10 ml	Hal wakhti
13-24 kg (28-54 rodol)	15 ml	Hal wakhti