

Warning!

If you are taking any medicines on this list, you need to **call 211 or your doctor right away.**

- Digoxin – a heart medicine
- Probenecid – for gout
- Diabetes medicines, insulin or pills
- Rifampin – for tuberculosis
- Coumadin or warfarin – blood thinner
- Accutane – for acne
- Theophylline – a breathing medicine
- Dilantin, barbiturates, or carbamazepine – seizure medicines
- Lithium – for mental health
- Cyclosporine – for immune system problems
- Methotrexate – for arthritis, cancer, psoriasis or lupus
- Requip – for Parkinson’s disease

Taking the emergency antibiotics with any of these medicines may cause serious, even life threatening, side effects.

Your doctor may want to change the medicine or the amount you take, or the doctor may want you to have more tests.

警告！

如果您正在服用此清單上列出的藥物，您需要**立即打電話211或打電話給您的醫生。**

- Digoxin – 心臟藥物
- Probenecid – 治療痛風的藥物
- 糖尿病藥物，胰島素或藥丸
- Rifampin – 治療結核病的藥物
- Coumadin或warfarin – 血液稀釋藥
- Accutane – 痤瘡治療藥物
- Theophylline – 呼吸藥物
- Dilantin、barbiturates或carbamazepine – 治療癲癇的藥物
- Lithium – 精神藥物
- Cyclosporine – 免疫系統藥物
- Methotrexate – 關節炎、癌症、銀屑病或狼瘡藥物
- Requip – 巴金森病藥物

如果上述任何藥物與日常應急用的抗生素同時使用，可能導致嚴重或甚至有生命危險的副作用。

醫生可能會調整您使用的藥物種類或劑量，或要求您接受更多的測試。